
Comoros

2010 total population: 734 750 Income group: Low

NCD mortality*			
2008 estimates		males	females
Total NCD deaths (000s)		1.1	1.3
NCD deaths under age 60 (percent of all NCD deaths)		46.3	45.1
Age-standardized death rate per 100 000			
All NCDs		798.4	767.7
Cancers		87.9	93.2
Chronic respiratory diseases		116.0	71.5
Cardiovascular diseases and diabetes		433.3	467.8
Behavioural risk factors			
2008 estimated prevalence (%)	males	females	total
Current daily tobacco smoking	15.6	6.8	11.2
Physical inactivity	4.8	9.1	6.9
Metabolic risk factors			
2008 estimated prevalence (%)	males	females	total
Raised blood pressure			
Raised blood glucose			
Overweight	18.0	21.6	19.8
Obesity	3.2	5.5	4.4
Raised cholesterol			




NCDs are estimated to account for 40% of all deaths.

No

0/5

Metabolic risk factor trends



NCD morbidity Yes Tobacco NCD risk factors No Number of tobacco (m)POWER measures Has a national, population-based cancer registry No implemented at the highest level of achievement

* The mortality estimates for this country have a high degree of uncertainty because they are not based on any national NCD mortality data. The estimates are based on a combination of country life tables cause of death models, regional cause of death patterns, and WHO and UNAIDS program estimates for some major causes of death (not including NCDs).

... = no data available

